



Telefon: 49(0) 4721 78 0
49(0) 4721 78 5 emergency call
 Fax: 49(0) 4721 78 1520
 E-mail: medico@tmas-germany.de
 Internet: www.tmas-germany.de
 MRCC Bremen: 49(0) 421 536870

Anleitung zur Gesundheitspflege Auf Kauffahrteischiffen

Chapter Attach. B No.

6. Stomach Ache

Possible causes	B.1.3.2. B.3.1.7. B.3.1.2.3. B.3.1.3.3. B.3.1.3.7. B.3.1.3.9. B.3.1.3.8. B.3.1.5.3.1. B.3.1.5. B.3.2.1.2.1. B.3.4.1.4. B.3.7.3.1. B.2.1.2.7.1 u. B.2.1.2.7.2.	
<ul style="list-style-type: none"> * "Acute stomach ache" <ul style="list-style-type: none"> caused by: stomach ulcer, with and without perforation appendicitis peritonitis intestinal occlusion strangulated hernia gallstone with colic inflammation of the gall bladder * Heart attack * Renal colic * Pneumonia and pleurisy in the area of the diaphragm * Accident with and without visible external injuries 		
<u>Rescue</u> <ul style="list-style-type: none"> * Illnesses in the digestive tract: Lay flat, prop up head and legs * Heart attack: Begin treatment before transporting. Administer oxygen pain treatment, vasodilator * Accident: Apply infusion before rescue measures, lay flat, prop up head and legs 	C.5. C.1.7.4. C.1.7.3. C.1.5. C.1.7.4.	25.01 23.02, 23.03, 2.02 11.01, 25.01
<u>First Aid</u> <ul style="list-style-type: none"> * Be prepared for heart and lung resuscitation * Comfortable positioning, back position, prop up knees and head * Have spit-basin and cellulose ready * Do not give anything to eat or drink * Pain treatment: depending on illness and medical instructions * Apply infusion 	C.1.1. C.1.7.4. C.1.5.	23.02 – 23.08, 25.02 0.13 167, 202 11.01
<u>Further Treatment</u> <p>Aim for diagnosis</p> <ul style="list-style-type: none"> * Peritonitis: After e.g. perforated stomach-intestinal ulcer, appendicitis: position flat, prop up head and legs, apply infusion, absolutely no intake of solids * Colic: With kidney stones: Drink a lot, walk around, painkiller Gall stones: painkiller. Nutrition: Leave out coffee, eggs, fried or grilled food * Strangulated hernia: Try to push it back in the lying position, with drawn up legs, no solid food * Heart attack: Influence in a calm way, painkiller, vasodilator, administer oxygen, absolute bed rest, * Pneumonia and pleurisy: bed rest, upper body high, hot chest compresses, painkiller, administer oxygen if required * Accident with and without visible external injuries: Infusions, painkiller, back splinting, administer oxygen if required, in the case of fractured ribs lay on injured side, pneumothorax ? 	B.1.3.2. B.3.1.3.7. C.1.7.4. C.1.5. B.3.4.1.4.-5/6 B3.1.5.3.1. B.3.1.3.8. B.3.2.1.2.1. B.3.7.3.1. B.2.1.2.7.1./2 B.2.1.2.2.2./3	5.06, 5.07, 8.02, 8.03 11.01 5.05, 5.06 5.05, 5.07, 2.02 5.06 2.01, 5.08, 6.02 23.02 8.01, 1.01, 5.02 23.01, 5.06, 5.08, 11.01, 23.01 5.06
<u>Further Measures:</u> <ul style="list-style-type: none"> * Frequent checking of respiration, consciousness, circulation (RCC-rule : <u>Respiration</u>, <u>Consciousness</u>, <u>Circulation</u>) * Make medical appointment * Telemetry of ECG, blood pressure, oxygenation, respiration * Following an accident, lay patient in vacuum mattress 		